arXiv:0811.3727v1 [math-ph] 23 Nov 2008

Algebraic Approaches to the Geopotential Forecast and Nonlinear MHD Equations[1]

Xiaoping Xu

Institute of Mathematics, Academy of Mathematics & System Sciences

Chinese Academy of Sciences, Beijing 100190, P.R. China [2]

Abstract

In this paper, we use various anstazes motivated from our earlier works on transonic gas flows, boundary layer problems and Navier-Stokes equations to find new explicit exact solutions with multiple parameter functions for the equation of geopotential forecast and the equations of nonlinear magnetohydrodynamics.

1 Introduction

Partial differential equation

$$(H_{xx}+H_{yy})_t+H_x(H_{xx}+H_{yy})_y-H_y(H_{xx}+H_{yy})_x=kH_x \tag{1.1}$$

is used in earth sciences for geopotential forecast on a middle level (e.g., cf [Ki] and Page 222 in [I]), where k is a real constant. Kibel' [Ki] found the Gaurvitz solution of the above equation. The well known Syono solution was given in [Sy]. Katkov [Ka1, Ka2] determined the Lie point symmetries and obtained certain invariant solutions. The other known solutions are related to the physical backgrounds such as configuration of type of narrow gullies and crests, flows of type of isolate whirlwinds, stream flow, springs and drains, hyperbolic points, and cyclone formation (e.g., cf. Pages 225, 226 in [I]).

In magnetohydrodynamics, it is very important to study the nonlinear MHD equations:

$$\psi_t+\varphi_x\psi_y-\varphi_y\psi_x=\varphi_z, \tag{1.2}$$

$$\begin{aligned}&(\varphi_{xx}+\varphi_{yy})_t+\varphi_x(\varphi_{xx}+\varphi_{yy})_y-\varphi_y(\varphi_{xx}+\varphi_{yy})_x\\=\;&(\psi_{xx}+\psi_{yy})_z+\psi_x(\psi_{xx}+\psi_{yy})_y-\psi_y(\psi_{xx}+\psi_{yy})_x,\end{aligned} \tag{1.3}$$

where φ and ψ are the potentials for the velocity and the transverse component of the magnetic field, respectively (they can also be interpreted as the potential of an electric field and the z-component of the vector potential of the magnetic field) (e.g., cf. [KP] and Page 390 in [I]). We refer [P] for more information on magnetohydrodynamics. Samokhin [S] (1985) determined the Lie point symmetries of the above equations, conservation laws and some solutions in terms the solutions of the other partial differential equations. Bershadskii [B] found a connection

---

[1] 2000 Mathematical Subject Classification. Primary 35C05, 35Q35; Secondary 35C10, 35C15.

[2] Research supported by China NSF 10871193

between the energy conservation law and the uniqueness of classical solution of the nonlinear MHD equations. The reason of solving the geopotential equation together with the nonlinear MHD equations is that the left hand side of the equation (1.1) coincides with that of the equation (1.3).

Based our earlier works on transonic gas flows [X1], boundary layer problems [X2] and Navier-Stokes equations [X3], we give in this paper various ansatzes related to algebraic characteristics of the above equations to find new explicit exact solutions with multiple parameter functions. By specifying these parameter functions, one can obtain the solutions of certain initial-value problems of the above equations.

The symmetry group of the geopotential equation (1.1) is generated by the following transformations:

$$T_{a,b}(H) = H(t+a,x,y+b), \qquad T_c(H) = c^3 H(c^{-1}t,cx,cy), \qquad a,b,c \in \mathbb{R},\ c \neq 0; \tag{1.4}$$

$$T_{\alpha,\beta}(H) = H(t,x+\alpha,y) + \alpha' y + \beta, \tag{1.5}$$

where α and β are arbitrary functions of t. The symmetry transformations of the nonlinear MHD equations (1.2) and (1.3) that we are concerned with are:

$$T_{a,b}(\varphi) = \varphi(t+a,x,y,z+b), \qquad T_{a,b}(\psi) = \psi(t+a,x,y,z+b), \qquad a,b \in \mathbb{R}; \tag{1.6}$$

$$T_{1c}(\varphi) = c^{-1}\varphi(ct,x,y,cz), \qquad T_{1c}(\psi) = c^{-1}\psi(ct,x,y,cz), \tag{1.7}$$

$$T_{2c}(\varphi) = c^{-2}\varphi(t,cx,xy,c), \qquad T_{2c}(\psi) = c^{-2}\psi(t,cx,cy,z) \tag{1.8}$$

for $0 \neq c \in \mathbb{R}$;

$$T_{\sigma,\tau}(\varphi) = \varphi(t,x+\sigma,y,z) + \sigma_t y + \tau_t, \qquad T_{\sigma,\tau}(\psi) = \psi(t,x+\sigma,y,z) + \sigma_z y + \tau_z, \tag{1.9}$$

$$T_{\sigma}(\varphi) = \varphi(t,x,y+\sigma,z) - \sigma_t x, \qquad T_{\sigma}(\psi) = \psi(t,x,y+\sigma,z) - \sigma_z x, \tag{1.10}$$

where σ and τ are any functions of t, z;

$$T_{\alpha}(\varphi) = \varphi(t,x\cos 2\alpha + y\sin 2\alpha, -x\sin 2\alpha + y\cos 2\alpha, z) + \alpha'(x^2+y^2), \tag{1.11}$$

$$T_{\alpha}(\psi) = \psi(t,x\cos 2\alpha + y\sin 2\alpha, -x\sin 2\alpha + y\cos 2\alpha, z) + \alpha'(x^2+y^2), \tag{1.12}$$

$$T_{\beta}(\varphi) = \varphi(t,x\cos 2\beta + y\sin 2\beta, -x\sin 2\beta + y\cos 2\beta, z) + \beta'(x^2+y^2), \tag{1.13}$$

$$T_{\beta}(\psi) = \psi(t,x\cos 2\beta + y\sin 2\beta, -x\sin 2\beta + y\cos 2\beta, z) - \beta'(x^2+y^2), \tag{1.16}$$

where $\alpha = \alpha(t+z)$ and $\beta = \beta(t-z)$ are arbitrary one-variable functions. The above transformations change solutions to solutions. For convenience, we always assume that all the involved partial derivatives of related functions always exist and we can change orders of taking partial derivatives. We also use prime $'$ to denote the derivative of any one-variable function.

In Section 2, we solve the equation (1.1) of geopotential forecast. We find explicit exact solutions of the nonlinear MHD equations (1.2) and (1.3) in Section 3.

2 Solutions of Geopotential Forecast Equation

In this section, we find two families of exact solutions of the geopotential forecast equation (1.1).

Let α and β be functions of t. Set

$$\varpi = \alpha x + \beta y. \tag{2.1}$$

Assume

$$H = \phi(t,\varpi) + \mu y^2 + \tau x + \nu y, \tag{2.2}$$

where ϕ is a two-variable function and τ, μ, ν are functions in t. Note

$$H_x = \alpha\phi_\varpi + \tau, \qquad H_y = \beta\phi_\varpi + 2\mu y + \nu, \qquad H_{xx} + H_{yy} = 2\mu + (\alpha^2 + \beta^2)\phi_{\varpi\varpi}, \tag{2.3}$$

$$(H_{xx} + H_{yy})_t = 2\mu' + (\alpha^2 + \beta^2)'\phi_{\varpi\varpi} + (\alpha^2 + \beta^2)[\phi_{t\varpi\varpi} + (\alpha' x + \beta' y)\phi_{\varpi\varpi\varpi}], \tag{2.4}$$

$$(H_{xx} + H_{yy})_x = (\alpha^2 + \beta^2)\alpha\phi_{\varpi\varpi\varpi}, \qquad (H_{xx} + H_{yy})_y = (\alpha^2 + \beta^2)\beta\phi_{\varpi\varpi\varpi}. \tag{2.5}$$

Thus (1.1) becomes

$$\begin{aligned} &2\mu' + (\alpha^2 + \beta^2)'\phi_{\varpi\varpi} + (\alpha^2 + \beta^2)\phi_{t\varpi\varpi} - k(\alpha\phi_\varpi + \tau) \\ &+(\alpha^2 + \beta^2)[\alpha' x + (\beta' - 2\alpha\mu)y + \beta\tau - \alpha\nu]\phi_{\varpi\varpi\varpi} = 0. \end{aligned} \tag{2.6}$$

In order to solve the above equation, we assume

$$2\mu' = k\tau, \qquad \tau = \alpha\vartheta'', \qquad \nu = \beta\vartheta'', \tag{2.7}$$

for some function ϑ of t, and

$$\alpha' x + (\beta' - 2\alpha\mu)y = 0. \tag{2.8}$$

Note that (2.8) is equivalent to the following system of ordinary differential equations:

$$\alpha' = 0, \qquad \beta' - 2\alpha\mu = 0. \tag{2.9}$$

By the first equation in (2.9), we have $\alpha = c \in \mathbb{R}$. So $\tau = c\vartheta''$ according to the second equation in (2.7). Moreover, the first equation in (2.7) yield

$$\mu = \frac{kc\vartheta' + c_0}{2}, \qquad c_0 \in \mathbb{R}. \tag{2.10}$$

Hence the second equation in (2.9) becomes

$$\beta' - c(kc\vartheta' + c_0) = 0. \tag{2.11}$$

Therefore,

$$\beta = c(kc\vartheta + c_0 t) + d, \qquad d \in \mathbb{R}. \tag{2.12}$$

According to the third equation in (2.7),

$$\nu = [c(kc\vartheta + c_0 t) + d]\vartheta''. \tag{2.13}$$

Now (2.6) becomes

$$(\alpha^2 + \beta^2)'\phi_{\varpi\varpi} + (\alpha^2 + \beta^2)\phi_{t\varpi\varpi} - kce^\gamma\phi_\varpi = 0. \tag{2.14}$$

Modulo the transformation in (1.5), it is enough to solve the following equation:

$$(\alpha^2 + \beta^2)'\phi_\varpi + (\alpha^2 + \beta^2)\phi_{t\varpi} - kce^\gamma\phi = 0. \tag{2.15}$$

The above equation can written as

$$[(\alpha^2 + \beta^2)\phi_\varpi]_t - kc\phi = 0. \tag{2.16}$$

So we take the form

$$\phi = \frac{\hat{\phi}(t, \varpi)}{\alpha^2 + \beta^2} = \frac{\hat{\phi}(t, \varpi)}{c^2 + [c(kc\vartheta + c_0 t) + d]^2}. \tag{2.17}$$

Then (2.16) becomes

$$\hat{\phi}_{\varpi t} = \frac{kc\hat{\phi}}{c^2 + [c(kc\vartheta + c_0 t) + d]^2}. \tag{2.18}$$

Thus we have the solution:

$$\begin{aligned} \hat{\phi} \quad = \quad & \sum_{i=1}^{m} d_i \exp\left(\frac{kca_i}{a_i^2 + b_i^2}\int \frac{dt}{c^2 + [c(kc\vartheta + c_0 t) + d]^2} + a_i\varpi\right) \\ & \times \sin\left(b_i\varpi + c_i - \frac{kcb_i}{a_i^2 + b_i^2}\int \frac{dt}{c^2 + d^2 e^{4kc^2\vartheta}}\right), \end{aligned} \tag{2.19}$$

where a_i, b_i, c_i, d_i are real constants such that $(a_i, b_i) \neq (0, 0)$.

Theorem 2.1. *Let ϑ be any function of t and let $a_i, b_i, c_i, d_i, c_0, c, d$ for $i = 1, ..., m$ be real constants such that $(c, d), (a_i, b_i) \neq (0, 0)$. We have the following solution of the geopotential forecast equation (1.1):*

$$\begin{aligned} H \quad = \quad & \frac{kc\vartheta' + c_0}{2}y^2 + \vartheta''[cx + [c(kc\vartheta + c_0 t) + d]y] + \frac{1}{c^2 + [c(kc\vartheta + c_0 t) + d]^2} \\ & \times \sum_{i=1}^{m} d_i \exp\left(\frac{kca_i}{a_i^2 + b_i^2}\int \frac{dt}{c^2 + [c(kc\vartheta + c_0 t) + d]^2} + a_i[cx + [c(kc\vartheta + c_0 t) + d]y]\right) \\ & \times \sin\left(b_i[cx + [c(kc\vartheta + c_0 t) + d]y] + c_i - \frac{kcb_i}{a_i^2 + b_i^2}\int \frac{dt}{c^2 + [c(kc\vartheta + c_0 t) + d]^2}\right). \end{aligned} \tag{2.20}$$

Next we set

$$\varpi = x^2 + y^2. \tag{2.21}$$

Assume

$$H = \xi(\varpi) - y \tag{2.22}$$

where ξ is a one-variable function. Note

$$H_x = 2x\xi', \qquad H_y = 2y\xi' - 1, \qquad H_{xx} + H_{yy} = 4(\xi' + \varpi\xi''), \tag{2.23}$$

$$(H_{xx} + H_{yy})_x = 8x(2\xi'' + \varpi\xi'''), \qquad (H_{xx} + H_{yy})_y = 8y(2\xi'' + \varpi\xi'''). \tag{2.24}$$

Then (1.1) is equivalent to:

$$4(2\xi'' + \varpi\xi''') = k\xi'. \tag{2.25}$$

Modulo the transformation (1.5), we only need to solve the equation:

$$\xi' + \varpi\xi'' = \frac{k}{4}\xi. \tag{2.26}$$

To solve the above ordinary differential equation, we assume

$$\xi = \sum_{i=0}^{\infty} \varpi^i(a_i + b_i \ln\varpi), \qquad a_i, b_i \in \mathbb{R}. \tag{2.27}$$

Observe

$$\xi' = \sum_{i=0}^{\infty} \varpi^{i-1}(ia_i + b_i + ib_i \ln\varpi), \tag{2.28}$$

$$\xi'' = \sum_{i=0}^{\infty} \varpi^{i-2}(i(i-1)a_i + (2i-1)b_i + i(i-1)b_i \ln\varpi). \tag{2.29}$$

So (2.26) becomes

$$\sum_{i=0}^{\infty}\varpi^{i-1}(i^2a_i+2ib_i+i^2b_i\ln\varpi)=\frac{k}{4}\sum_{i=0}^{\infty}\varpi^i(a_i+b_i\ln\varpi), \tag{2.30}$$

equivalently,

$$(i+1)^2a_{i+1}+2(i+1)b_{i+1}=\frac{k}{4}a_i, \qquad (i+1)^2b_{i+1}=\frac{k}{4}b_i. \tag{2.31}$$

Hence

$$b_i=\frac{b_0k^i}{(i!)^24^i}, \qquad a_i=\frac{a_0k^i}{(i!)^24^i}-2b_0\sum_{r=0}^{i}\frac{k^r}{r(r!)^24^r}. \tag{2.32}$$

Thus

$$\xi=a_0\sum_{i=0}^{\infty}\frac{k^i\varpi^i}{(i!)^24^i}+b_0\sum_{j=0}^{\infty}\varpi^j\left(\frac{k^j\ln\varpi}{(j!)^24^j}-2\sum_{r=0}^{j}\frac{k^r}{r(r!)^24^r}\right). \tag{2.33}$$

Theorem 2.2. *Let b and c be any real constants. We have the following steady solution of the geopotential forecast equation (1.1):*

$$H=-y+b\sum_{i=0}^{\infty}\frac{k^i(x^2+y^2)^i}{(i!)^24^i}+c\sum_{j=0}^{\infty}(x^2+y^2)^j\left(\frac{k^j\ln(x^2+y^2)}{(j!)^24^j}-2\sum_{r=0}^{j}\frac{k^r}{r(r!)^24^r}\right). \tag{2.34}$$

Remark 2.3. Applying the transformation (1.5), we obtain the following non-steady solution:

$$\begin{aligned}H &= (\alpha'-1)y+\beta+b\sum_{i=0}^{\infty}\frac{k^i((x+\alpha)^2+y^2)^i}{(i!)^24^i}\\ &\quad +c\sum_{j=0}^{\infty}((x+\alpha)^2+y^2)^j\left(\frac{k^j\ln((x+\alpha)^2+y^2)}{(j!)^24^j}-2\sum_{r=0}^{j}\frac{k^r}{r(r!)^24^r}\right),\end{aligned} \tag{2.35}$$

where α and β are arbitrary functions of t.

3 Solutions of Nonlinear MHD Equations

In this section, we will find multiple parameter function exact solutions of the nonlinear MHD equations (1.2) and (1.3).

We first assume

$$\varphi=\sigma_txy+fx+gy+h, \qquad \psi=\sigma_zxy+\lambda x+\mu y+\rho, \tag{3.1}$$

where f,g,h,σ,λ,μ and ρ are functions in t,z. Then (1.3) naturally holds. Moreover, (1.2) becomes

$$(\lambda_t-f_z)x+(\mu_t-g_z)y+\rho_t-h_z+(\sigma_ty+f)(\sigma_zx+\mu)-(\sigma_tx+g)(\sigma_zy+\lambda)=0, \tag{3.2}$$

equivalently,

$$\lambda_t-f_z+\sigma_zf-\sigma_t\lambda=0, \tag{3.3}$$

$$\mu_t-g_z+\sigma_t\mu-\sigma_zg=0, \tag{3.4}$$

$$\rho_t-h_z+f\mu-g\lambda=0. \tag{3.5}$$

Solving (3.3) and (3.4), we get

$$f = \vartheta_t e^{\sigma}, \qquad g = \tau_t e^{-\sigma}, \qquad \lambda = \vartheta_z e^{\sigma}, \qquad \mu = \tau_z e^{-\sigma}, \tag{3.6}$$

where ϑ and τ are arbitrary functions in t, z. Moreover, (3.5) becomes

$$\rho_t - h_z + \vartheta_t \tau_z - \vartheta_z \tau_t = 0. \tag{3.7}$$

Thus

$$h = \vartheta_t \tau, \qquad \rho = \vartheta_z \tau \tag{3.8}$$

modulo the transformation of type $T_{0,\nu}$ in (1.9). So we have the following simple conclusion.

Proposition 3.1. *Let σ, ϑ and τ be functions of t, z. We get the following solution of the nonlinear MHD equations (1.2) and (1.3):*

$$\varphi = \sigma_t xy + \vartheta_t e^{\sigma} x + \tau_t e^{-\sigma} y + \vartheta_t \tau, \tag{3.9}$$

$$\psi = \sigma_z xy + \vartheta_z e^{\sigma} x + \tau_z e^{-\sigma} y + \vartheta_z \tau. \tag{3.10}$$

The main objective in this section is to find more sophisticated exact solutions of the equations (1.2) and (1.3). Let $\Im$ and ε be functions in t, z. Set

$$\varpi = \Im x + \varepsilon y. \tag{3.11}$$

Suppose

$$\varphi = \zeta(t, z, \varpi) + \sigma_t xy, \qquad \psi = \eta(t, z, \varpi) + \sigma_z xy, \tag{3.12}$$

where ζ and η are functions in t, z, ϖ to be determined, and σ is a function of t, z. Then (1.2) becomes

$$\eta_t - \zeta_z + [\Im_t x + \varepsilon_t y - \sigma_t(\Im x - \varepsilon y)]\eta_{\varpi} - [\Im_z x + \varepsilon_z y - \sigma_z(\Im x - \varepsilon y)]\zeta_{\varpi} = 0 \tag{3.13}$$

and (1.3) becomes

$$\begin{aligned}&(\Im^2 + \varepsilon^2)_t \zeta_{\varpi\varpi} - (\Im^2 + \varepsilon^2)_z \eta_{\varpi\varpi} + (\Im^2 + \varepsilon^2)\{\zeta_{t\varpi\varpi} + [\Im_t x + \varepsilon_t y - \sigma_t(\Im x - \varepsilon y)]\zeta_{\varpi\varpi\varpi}\\ &-\eta_{z\varpi\varpi} - [\Im_z x + \varepsilon_z y - \sigma_z(\Im x - \varepsilon y)]\eta_{\varpi\varpi\varpi}\} = 0.\end{aligned} \tag{3.14}$$

In order to solve the above system of partial differential equation, we assume

$$\Im_s x + \varepsilon_s y - \sigma_s(\Im x - \varepsilon y) = 0, \qquad s = t, z, \tag{3.15}$$

equivalently,

$$\Im_s - \sigma_s \Im = 0, \qquad \varepsilon_s + \sigma_s \varepsilon = 0, \qquad s = t, z. \tag{3.16}$$

So

$$\Im = b e^{\sigma}, \qquad \varepsilon = c e^{-\sigma}, \qquad b, c \in \mathbb{R}. \tag{3.17}$$

Moreover, (3.13) becomes

$$\eta_t - \zeta_z = 0. \tag{3.18}$$

Hence

$$\zeta = F_t(t, z, \varpi), \qquad \eta = F_z(t, z, \varpi) \tag{3.19}$$

for some three variable function F. Now (3.14) becomes:

$$(\Im^2 + \varepsilon^2)_t F_{t\varpi\varpi} - (\Im^2 + \varepsilon^2)_z F_{z\varpi\varpi} + (\Im^2 + \varepsilon^2)(F_{tt\varpi\varpi} - F_{zz\varpi\varpi}) = 0. \tag{3.20}$$

Modulo the transformation in (1.9), we have

$$F = \hat{F} + \lambda\varpi \tag{3.21}$$

for a function λ of t, z and a function $\hat{F}$ of t, z, ϖ such that

$$(\Im^2 + \varepsilon^2)_t \hat{F}_t - (\Im^2 + \varepsilon^2)_z \hat{F}_z + (\Im^2 + \varepsilon^2)(\hat{F}_{tt} - \hat{F}_{zz}) = 0. \tag{3.22}$$

Rewrite (3.22) as

$$((\Im^2 + \varepsilon^2)\hat{F}_t)_t - ((\Im^2 + \varepsilon^2)\hat{F}_z)_z = 0. \tag{3.23}$$

In order to find exact solutions of the above equation, we take the following special cases of $\Im^2 + \varepsilon^2$. Without loss of generality, we assume $b = 1$ in (3.17).

Case 1. $\Im^2 + \varepsilon^2 = e^{a_1 t + a_2 z}$ with $a_1, a_2 \in \mathbb{R}$.

Our assumptions says

$$e^{2\sigma} + c^2 e^{-2\sigma} = e^{a_1 t + a_2 z}, \tag{3.24}$$

equivalently,

$$(e^{2\sigma})^2 - e^{a_1 t + a_2 z} e^{2\sigma} + c = 0. \tag{3.25}$$

So

$$e^{2\sigma} = \frac{e^{a_1 t + a_2 z} + \sqrt{e^{2(a_1 t + a_2 z)} - 4c^2}}{2} \tag{3.26}$$

or

$$e^{2\sigma} = \frac{e^{a_1 t + a_2 z} - \sqrt{e^{2(a_1 t + a_2 z)} - 4c^2}}{2}. \tag{3.27}$$

Hence

$$\sigma = \frac{1}{2}\left(\ln\left(e^{a_1 t + a_2 z} + \sqrt{e^{2(a_1 t + a_2 z)} - 4c^2}\right) - \ln 2\right) \tag{3.28}$$

or

$$\sigma = \frac{1}{2}\left(\ln\left(e^{a_1 t + a_2 z} - \sqrt{e^{2(a_1 t + a_2 z)} - 4c^2}\right) - \ln 2\right). \tag{3.29}$$

Now (3.23) is equivalent to

$$\hat{F}_{tt} + a_1 \hat{F}_t = \hat{F}_{zz} + a_2 \hat{F}_z. \tag{3.30}$$

For $0 \neq a \in \mathbb{R}$, we denote

$$\mathcal{D}(a, s) = a\partial_s + \partial_s^2 \tag{3.31}$$

and

$$\xi_{1,0}(a, s) = 1, \qquad \xi_{1,i}(a, s) = \sum_{r=0}^{i-1} \frac{(-1)^r s^{i-r}}{(i-r)! a^{i+r}}, \tag{3.32}$$

$$\xi_{2,0}(a, s) = e^{-as}, \qquad \xi_{2,i}(a, s) = (-1)^i e^{-as} \sum_{r=0}^{i-1} \frac{s^{i-r}}{(i-r)! a^{i+r}}, \tag{3.33}$$

for $0 < i \in \mathbb{Z}$. Moreover, we let

$$\mathcal{D}(0, s) = \partial_s^2, \qquad \xi_{1,i}(0, s) = \frac{s^{2i}}{(2s)!}, \qquad \xi_{2,i}(0, s) = \frac{s^{2i+1}}{(2i+1)!} \tag{3.34}$$

for $0 \leq i \in \mathbb{Z}$. Then

$$\mathcal{D}(a,s)(\xi_{\epsilon,0}(a,s)) = 0, \qquad \mathcal{D}(a,s)(\xi_{\epsilon,i}(a,s)) = \xi_{\epsilon,i-1}(a,s), \qquad \epsilon = 1,2, \tag{3.35}$$

for $0 < i \in \mathbb{Z}$. Given one-variable functions

$$\{\alpha_{\epsilon,i_\epsilon}(\varpi), \beta_{\epsilon,j_\epsilon}(\varpi) \mid \epsilon = 1,2;\ i_\epsilon = 0,1,...,m_\epsilon;\ j_\epsilon = 0,1,...,n_\epsilon\}, \tag{3.36}$$

we have the following solution of (3.30):

$$\hat{F} = \sum_{\epsilon=1,2}[\sum_{i=0}^{m_\epsilon} \alpha_{\epsilon,i}(\varpi)\xi_{\epsilon,i}(a_1,t)\xi_{1,m_\epsilon-i}(a_2,z) + \sum_{j=0}^{n_\epsilon} \beta_{\epsilon,j}(\varpi)\xi_{\epsilon,j}(a_1,t)\xi_{2,n_\epsilon-j}(a_2,z)]. \tag{3.37}$$

Case 2. Case 1. $\Im^2 + \varepsilon^2 = e^{a_1 t} z^{a_2}$ with $a_1, a_2 \in \mathbb{R}$ such that $a_2 \notin \{0, \mathbb{Z} + 1/2\}$.

As (3.24)-(3.28), we have:

$$\sigma = \frac{1}{2}\left(\ln\left(e^{a_1 t} z^{a_2} + \sqrt{e^{2a_1 t} z^{2a_2} - 4c^2}\right) - \ln 2\right) \tag{3.38}$$

or

$$\sigma = \frac{1}{2}\left(\ln\left(e^{a_1 t} z^{a_2} - \sqrt{e^{2a_1 t} z^{2a_2} - 4c^2}\right) - \ln 2\right). \tag{3.39}$$

In this case, (3.23) is equivalent to

$$\hat{F}_{tt} + a_1 \hat{F}_t = \hat{F}_{zz} + a_2 z^{-1} \hat{F}_z. \tag{3.40}$$

For $a \in \mathbb{R} \setminus \{0, \mathbb{Z} + 1/2\}$,

$$\zeta_{1,0}(a,s) = 1, \qquad \zeta_{1,i}(a,s) = \frac{s^{2i}}{2^i i! \prod_{r=1}^{i}(a + 2r - 1)}, \tag{3.41}$$

$$\zeta_{2,0}(a,s) = z^{1-a}, \qquad \zeta_{2,i}(a,s) = \frac{s^{2i+1-a}}{2^i i! \prod_{r=1}^{i}(2r + 1 - a)} \tag{3.42}$$

for $0 < i \in \mathbb{Z}$. Denote

$$\hat{\mathcal{D}}(a,s) = \partial_s^2 + \frac{a}{s}\partial_s. \tag{3.43}$$

Then

$$\hat{\mathcal{D}}(a,s)(\zeta_{\epsilon,0}(a,s)) = 0, \qquad \hat{\mathcal{D}}(a,s)(\zeta_{\epsilon,i}(a,s)) = \zeta_{\epsilon,i-1}(a,s), \qquad \epsilon = 1,2, \tag{3.44}$$

for $0 < i \in \mathbb{Z}$. Given the one-variable functions in (3.36), we have the following solution of (3.40):

$$\hat{F} = \sum_{\epsilon=1,2}[\sum_{i=0}^{m_\epsilon} \alpha_{\epsilon,i}(\varpi)\xi_{\epsilon,i}(a_1,t)\zeta_{1,m_\epsilon-i}(a_2,z) + \sum_{j=0}^{n_\epsilon} \beta_{\epsilon,j}(\varpi)\xi_{\epsilon,j}(a_1,t)\zeta_{2,n_\epsilon-j}(a_2,z)]. \tag{3.45}$$

Case 3. Case 1. $\Im^2 + \varepsilon^2 = t^{a_1} z^{a_2}$ with $a_1, a_2 \in \mathbb{R} \setminus \{0, \mathbb{Z} + 1/2\}$.

As (3.24)-(3.28), we have:

$$\sigma = \frac{1}{2}\left(\ln\left(t^{a_1} z^{a_2} + \sqrt{t^{2a_1} z^{2a_2} - 4c^2}\right) - \ln 2\right) \tag{3.46}$$

or

$$\sigma = \frac{1}{2}\left(\ln\left(t^{a_1} z^{a_2} - \sqrt{t^{2a_1} z^{2a_2} - 4c^2}\right) - \ln 2\right). \tag{3.47}$$

In this case, (3.23) is equivalent to

$$\hat{F}_{tt}+a_1t^{-1}\hat{F}_t=\hat{F}_{zz}+a_2z^{-1}\hat{F}_z. \tag{3.48}$$

Given the one-variable functions in (3.36), we have the following solution of (3.48):

$$\hat{F}=\sum_{\epsilon=1,2}[\sum_{i=0}^{m_\epsilon}\alpha_{\epsilon,i}(\varpi)\zeta_{\epsilon,i}(a_1,t)\zeta_{1,m_\epsilon-i}(a_2,z)+\sum_{j=0}^{n_\epsilon}\beta_{\epsilon,j}(\varpi)\zeta_{\epsilon,j}(a_1,t)\zeta_{2,n_\epsilon-j}(a_2,z)]. \tag{3.49}$$

In summary, we have the following theorem:

Theorem 3.2. *Let λ be any function of t,z and let c be arbitrary real constant. Suppose the one-variable functions in (3.36) are given. We have the following solution of the nonlinear MHD equations (1.2) and (1.3):*

$$\varphi=\sigma_txy+\lambda_t(e^\sigma x+ce^{-\sigma}y)+\hat{F}_t(t,z,e^\sigma x+ce^{-\sigma}), \tag{3.50}$$

$$\psi=\sigma_zxy+\lambda_z(e^\sigma x+ce^{-\sigma}y)+\hat{F}_z(t,z,e^\sigma x+ce^{-\sigma}), \tag{3.51}$$

where (1) σ is given in (3.28) or (3.29) with $a_1,a_2\in\mathbb{R}$ and $\hat{F}(t,z,\varpi)$ is given in (3.37) via (3.32)-(3.34); (2) σ is given in (3.38) or (3.39) with $a_1,a_2\in\mathbb{R}$ such that $a_2\not\in\{0,\mathbb{Z}+1/2\}$ and $\hat{F}(t,z,\varpi)$ is given in (3.44) via (3.32)-(3.34) and (3.41)-(3.42); (3) σ is given in (3.46) or (3.47) with $a_1,a_2\in\mathbb{R}\setminus\in\{0,\mathbb{Z}+1/2\}$ and $\hat{F}(t,z,\varpi)$ is given in (3.49) via (3.41) and (3.42).

Finally, we have the following obvious results.

Proposition 3.3. *Let $F(w,\varpi)$ and $G(w,\varpi)$ be any two-variable functions. We have the following solutions of the nonlinear MHD equations (1.2) and (1.3):*

$$(1)\qquad \varphi=F_w(t+z,x)+G_w(t-z,x),\qquad \psi=F_w(t+z,x)-G_w(t-z,x); \tag{3.52}$$

(2)

$$\varphi=F_w(t+z,x^2+y^2)+G_w(t-z,x^2+y^2), \tag{3.53}$$

$$\psi=F_w(t+z,x^2+y^2)-G_w(t-z,x^2+y^2). \tag{3.54}$$

We remark that we would get more sophisticated solutions if we apply the transformations (1.6)-(1.16) to our above solutions. For instance, applying the transformations (1.11)-(1.16) to the solution in (3.52), we obtain the following solution of the nonlinear MHD equations (1.2) and (1.3):

$$\begin{aligned}\varphi &= F_w(t+z,(x+\sigma)\cos 2\alpha+(y+\tau)\sin 2\alpha)+\alpha'((x+\sigma)^2+(y+\tau)^2)\\ &\quad +\sigma_t(y+\tau)-\tau_tx+\lambda_t+G_w(t-z,(x+\sigma)\cos 2\alpha+(y+\tau)\sin 2\alpha),\end{aligned} \tag{3.55}$$

$$\begin{aligned}\psi &= F_w(t+z,(x+\sigma)\cos 2\alpha+(y+\tau)\sin 2\alpha)+\epsilon\alpha'((x+\sigma)^2+(y+\tau)^2)\\ &\quad +\sigma_t(y+\tau)-\tau_tx+\lambda_t-G_w(t-z,(x+\sigma)\cos 2\alpha+(y+\tau)\sin 2\alpha)\end{aligned} \tag{3.56}$$

with $\epsilon=\pm1$, where $\alpha=\alpha(t+\epsilon z)$ is an arbitrary one-variable function, and σ,τ,λ are any functions in t,z.

References

[B] A. G. Bershadskii, Energy conservation law and uniqueness of the classical solutions of nonlinear MHD problems, *Magnetohydrodynamics* **2** (1980), 78-80.

[I] N. H. Ibragimov, *Lie Group Analysis of Differential Equations*, Volume 2, CRC Handbook, CRC Press, 1995.

[LRT] C. C. Lin, E. Reissner and H. S. Tsien, On two-dimensional non-steady motion of a slender body in a compressible fluid, *J. Math. Phys.* **27** (1948), no. 3, 220.

[Ka1] V. I. Katkov, One class of exact solutions of the geopotential forecast equation, *Izvestiya Akad. Nauk S.S.S.R., Fizica Atmosferi i Okcana* **1** (1965), 1088.

[Ka2] V. I. Katkov, Exact solutions of the geopotential forecast equation, *Izvestiya Akad. Nauk S.S.S.R., Fizica Atmosferi i Okcana* **2** (1966), 1193.

[Ki] T. W. Kibel', *Introduction to Hydrodynamical Methods of Short Term Weather Forecast*, Gosteoretizdat, Moscow, 1954.

[M] E. V. Mamontov, On the theory of nonstationary transonic flows, *Dokl. Acad. Nauk SSSR* **185** (1969), no. 3, 538.

[O] L. V. Ovsiannikov, *Group Analysis of Differential Equations*, Academic Press, New York, 1982.

[P] Shih-I Pai, *Magnetogasdynamic and Plasma Physics*, Springer-Verlag, Wien, 1962.

[Sa] A. V. Samokhin, Nonlinear MHD equations: symmetries, solutions, and conservation laws, *Dokl. Akad, Nauk S.S.S.R.* **285** (1985), no. 5, 1101-1106.

[Sy] S. Syono, Various simple solutions of the barotropic vorticity equation, in *Vortex, Collected papers of the numerical weather prediction group in Tokyo* (1958), 3.

[X1] X. Xu, Stable-Range approach to the equation of nonstationary transonic gas flows, *Quart. Appl. Math.* **65** (2007), 529-547.

[X2] X. Xu, New algebraic approaches to classical boundary layer problems, *arXiv:0706.1864*.

[X3] X. Xu, Asymmetric and moving-frame approaches to Navier-Stokes equations, *Quart. Appl. Math.*, in press, arXiv:0706.1861.